

Nonlinear Control

Lecture # 28

Robust State Feedback Stabilization

Sliding Mode Control

$$\dot{x} = f(x) + B(x)[G(x)u + \delta(t, x, u)]$$

$x \in R^n$, $u \in R^m$, f and B are known, while G and δ could be uncertain, $f(0) = 0$, $G(x)$ is a positive definite symmetric matrix with

$$\lambda_{\min}(G(x)) \geq \lambda_0 > 0$$

Regular Form:

$$\begin{bmatrix} \eta \\ \xi \end{bmatrix} = T(x), \quad \frac{\partial T}{\partial x} B(x) = \begin{bmatrix} 0 \\ I \end{bmatrix}$$

$$\dot{\eta} = f_a(\eta, \xi), \quad \dot{\xi} = f_b(\eta, \xi) + G(x)u + \delta(t, x, u)$$

$$\dot{\eta} = f_a(\eta, \xi), \quad \dot{\xi} = f_b(\eta, \xi) + G(x)u + \delta(t, x, u)$$

Sliding Manifold:

$$s = \xi - \phi(\eta) = 0, \quad \phi(0) = 0$$

$$s(t) \equiv 0 \Rightarrow \dot{\eta} = f_a(\eta, \phi(\eta))$$

Design ϕ s.t. the origin of $\dot{\eta} = f_a(\eta, \phi(\eta))$ is asymp. stable

$$\dot{s} = f_b(\eta, \xi) - \frac{\partial \phi}{\partial \eta} f_a(\eta, \xi) + G(x)u + \delta(t, x, u)$$

$$u = \psi(\eta, \xi) + v$$

Typical choices of ψ :

$$\psi = 0, \quad \psi = -\hat{G}^{-1}[f_b - (\partial \phi / \partial \eta) f_a]$$

$$\dot{s} = G(x)v + \Delta(t, x, v)$$

$$\left\| \frac{\Delta(t, x, v)}{\lambda_{\min}(G(x))} \right\| \leq \varrho(x) + \kappa_0 \|v\|, \quad \forall (t, x, v) \in [0, \infty) \times D \times R^m$$

$$\varrho(x) \geq 0, \quad 0 \leq \kappa_0 < 1 \quad (\textit{Known})$$

$$V = \frac{1}{2}s^T s \Rightarrow \dot{V} = s^T \dot{s} = s^T G(x)v + s^T \Delta(t, x, v)$$

$$v = -\beta(x) \frac{s}{\|s\|}, \quad \beta(x) \geq \frac{\varrho(x)}{1 - \kappa_0} + \beta_0, \quad \beta_0 > 0$$

$$\begin{aligned} \dot{V} &= -\beta(x)s^T G(x)s/\|s\| + s^T \Delta(t, x, v) \\ &\leq \lambda_{\min}(G(x))[-\beta(x) + \varrho(x) + \kappa_0\beta(x)] \|s\| \\ &= \lambda_{\min}(G(x))[-(1 - \kappa_0)\beta(x) + \varrho(x)] \|s\| \\ &\leq -\lambda_{\min}(G(x))\beta_0(1 - \kappa_0)\|s\| \\ &\leq -\lambda_0\beta_0(1 - \kappa_0)\|s\| = -\lambda_0\beta_0(1 - \kappa_0)\sqrt{2V} \end{aligned}$$

Trajectories reach the manifold $s = 0$ in finite time and cannot leave it

Continuous Implementation

$$\text{Sat}(y) = \begin{cases} y, & \text{if } \|y\| \leq 1 \\ y/\|y\|, & \text{if } \|y\| > 1 \end{cases}$$

$$v = -\beta(x) \text{Sat}\left(\frac{s}{\mu}\right)$$

$$\|s\| \geq \mu \Rightarrow \text{Sat}(s/\mu) = s/\|s\| \Rightarrow s^T \dot{s} \leq -\lambda_0 \beta_0 (1 - \kappa_0) \|s\|$$

Trajectories reach the boundary layer $\{\|s\| \leq \mu\}$ in finite time and remains inside thereafter

Study the behavior of η : $\dot{\eta} = f_a(\eta, \phi(\eta) + s)$

$$\alpha_1(\|\eta\|) \leq V_0(\eta) \leq \alpha_2(\|\eta\|)$$

$$\frac{\partial V_0}{\partial \eta} f_a(\eta, \phi(\eta) + s) \leq -\alpha_3(\|\eta\|), \quad \forall \|\eta\| \geq \alpha_4(\|s\|)$$

$$\|s\| \leq c \Rightarrow \dot{V}_0 \leq -\alpha_3(\|\eta\|), \quad \text{for } \|\eta\| \geq \alpha_4(c)$$

$$\alpha(r) = \alpha_2(\alpha_4(r))$$


$$\begin{aligned} V_0(\eta) \geq \alpha(c) &\Leftrightarrow V_0(\eta) \geq \alpha_2(\alpha_4(c)) \Rightarrow \alpha_2(\|\eta\|) \geq \alpha_2(\alpha_4(c)) \\ &\Rightarrow \|\eta\| \geq \alpha_4(c) \\ &\Rightarrow \dot{V}_0 \leq -\alpha_3(\|\eta\|) \leq -\alpha_3(\alpha_4(c)) \end{aligned}$$

$$\Omega = \{V_0(\eta) \leq c_0\} \times \{\|s\| \leq c\}, \quad c_0 \geq \alpha(c), \quad \Omega \subset T(D)$$

$$V_0(\eta) \geq \alpha(\mu) \Rightarrow \dot{V}_0 \leq -\alpha_3(\alpha_4(\mu))$$

$\Rightarrow \Omega_\mu = \{V_0(\eta) \leq \alpha(\mu)\} \times \{\|s\| \leq \mu\}$ is positively invariant

In summary, all trajectories starting in Ω remain in Ω and reach Ω_μ in finite time and remain inside thereafter



Theorem 10.1

Suppose all the assumptions hold over Ω . Then, for all $(\eta(0), \xi(0)) \in \Omega$, the trajectory $(\eta(t), \xi(t))$ is bounded for all $t \geq 0$ and reaches the positively invariant set Ω_μ in finite time. If the assumptions hold globally and $V(\eta)$ is radially unbounded, the foregoing conclusion holds for any initial state

Example 10.2 (Magnetic levitation - friction neglected)

$$\dot{x}_1 = x_2, \quad \dot{x}_2 = 1 + \frac{m_o}{m}u, \quad x_1 \geq 0, \quad -2 \leq u \leq 0$$

We want to stabilize the system at $x_1 = 1$. Nominal steady-state control is $u_{ss} = -1$

Shift the equilibrium point to the origin: $x_1 \rightarrow x_1 - 1, u \rightarrow u + 1$

$$\dot{x}_1 = x_2, \quad \dot{x}_2 = \frac{m - m_o}{m} + \frac{m_o}{m}u$$

$$x_1 \geq -1, \quad |u| \leq 1$$

$$\text{Assume} \quad \left| \frac{(m - m_o)}{m_o} \right| \leq \frac{1}{3}$$

$$s = x_1 + x_2 \quad \Rightarrow \quad \dot{x}_1 = -x_1 + s$$

$$V_0 = \frac{1}{2}x_1^2$$

$$\dot{V}_0 = -x_1^2 + x_1 s \leq -(1-\theta)x_1^2, \quad \forall |x_1| \geq |s|/\theta, \quad 0 < \theta < 1$$

$$\alpha_1(r) = \alpha_2(r) = \frac{1}{2}r^2, \quad \alpha_3(r) = (1-\theta)r^2, \quad \alpha_4(r) = r/\theta$$

$$\alpha(r) = \alpha_2(\alpha_4(r)) = \frac{1}{2}(r/\theta)^2$$

$$\text{With } c_0 = \alpha(c), \quad \Omega = \{|x_1| \leq c/\theta\} \times \{|s| \leq c\}$$

$$\Omega_\mu = \{|x_1| \leq \mu/\theta\} \times \{|s| \leq \mu\}$$

$$\Omega = \{|x_1| \leq c/\theta\} \times \{|s| \leq c\}$$

Take $c \leq \theta$ to meet the constraint $x_1 \geq -1$

$$\dot{s} = x_2 + \frac{m - m_o}{m} + \frac{m_o}{m}u$$

$$\left| \frac{x_2 + (m - m_o)/m}{m_o/m} \right| = \left| \frac{m}{m_o}x_2 + \frac{m - m_o}{m_o} \right| \leq \frac{1}{3}(4|x_2| + 1)$$

$$\text{In } \Omega, \quad |x_2| \leq |x_1| + |s| \leq c(1 + 1/\theta)$$

$$\text{with } \frac{1}{\theta} = 1.1, \quad \left| \frac{x_2 + (m - m_o)/m}{m_o/m} \right| \leq \frac{8.4c + 1}{3}$$

To meet the constraint $|u| \leq 1$ limit c to

$$\frac{8.4c + 1}{3} < 1 \Leftrightarrow c < 0.238 \text{ and take } u = -\text{sat}\left(\frac{s}{\mu}\right)$$

With $c = 0.23$, Theorem 10.1 ensures that all trajectories starting in Ω stay in Ω and enter Ω_μ in finite time

$$\text{Inside } \Omega_\mu, \quad |x_1| \leq \mu/\theta = 1.1\mu$$

μ can be chosen small enough to meet any specified ultimate bound on x_1

$$\text{For } |x_1| \leq 0.01, \text{ take } \mu = 0.01/1.1 \approx 0.009$$

With further analysis inside Ω_μ we can derive a less conservative estimate of the ultimate bound of $|x_1|$. In Ω_μ , the closed-loop system is represented by

$$\dot{x}_1 = x_2, \quad \dot{x}_2 = \frac{m - m_o}{m} - \frac{m_o(x_1 + x_2)}{m\mu}$$

which has a unique equilibrium point at

$$\left(x_1 = \frac{\mu(m - m_o)}{m_o}, x_2 = 0 \right)$$

and its matrix is Hurwitz

$$\lim_{t \rightarrow \infty} x_1(t) = \frac{\mu(m - m_o)}{m_o}, \quad \lim_{t \rightarrow \infty} x_2(t) = 0$$

$$\left| \frac{(m - m_o)}{m_o} \right| \leq \frac{1}{3} \Rightarrow |x_1| \leq 0.34\mu$$

For $|x_1| \leq 0.01$, take $\mu = 0.029$

We can also obtain a less conservative estimate of the region of attraction

$$V_1 = \frac{1}{2}(x_1^2 + s^2)$$

$$\dot{V}_1 \leq -x_1^2 + s^2 - \frac{m_o}{m} \left[1 - \left| \frac{m - m_o}{m_o} \right| \right] |s| \leq -x_1^2 + s^2 - \frac{1}{2}|s|$$

for $|s| \geq \mu$

$$\dot{V}_1 \leq -x_1^2 + s^2 + \left| \frac{m - m_o}{m_o} \right| |s| - \frac{m_o}{m} \frac{s^2}{\mu} \leq -x_1^2 + s^2 + \frac{1}{2}|s| - \frac{3s^2}{4\mu}$$

for $|s| \leq \mu$

With $\mu = 0.029$, it can be verified that $\dot{V}_1$ is less than a negative number in the set $\{0.0012 \leq V_1 \leq 0.12\}$. Therefore, all trajectories starting in $\Omega_1 = \{V_1 \leq 0.12\}$ enter $\Omega_2 = \{V_1 \leq 0.0012\}$ in finite time. Since $\Omega_2 \subset \Omega$, our earlier analysis holds and the ultimate bound of $|x_1|$ is 0.01. The new estimate of the region of attraction, Ω_1 , is larger than Ω

