

Nonlinear Control

Lecture # 2

Stability of Equilibrium Points

Basic Concepts

$$\dot{x} = f(x)$$

f is locally Lipschitz over a domain $D \subset \mathbb{R}^n$

Suppose $\bar{x} \in D$ is an equilibrium point; that is, $f(\bar{x}) = 0$

Characterize and study the stability of $\bar{x}$

For convenience, we state all definitions and theorems for the case when the equilibrium point is at the origin of $\mathbb{R}^n$; that is, $\bar{x} = 0$. No loss of generality

$$y = x - \bar{x}$$

$$\dot{y} = \dot{x} = f(x) = f(y + \bar{x}) \stackrel{\text{def}}{=} g(y), \quad \text{where } g(0) = 0$$

Definition 3.1

The equilibrium point $x = 0$ of $\dot{x} = f(x)$ is

- stable if for each $\varepsilon > 0$ there is $\delta > 0$ (dependent on ε) such that

$$\|x(0)\| < \delta \Rightarrow \|x(t)\| < \varepsilon, \quad \forall t \geq 0$$


- unstable if it is not stable
- asymptotically stable if it is stable and δ can be chosen such that

$$\|x(0)\| < \delta \Rightarrow \lim_{t \rightarrow \infty} x(t) = 0$$


Scalar Systems ($n = 1$)

The behavior of $x(t)$ in the neighborhood of the origin can be determined by examining the sign of $f(x)$


The ε - δ requirement for stability is violated if $xf(x) > 0$ on either side of the origin



Unstable




Unstable




Unstable


The origin is stable if and only if $xf(x) \leq 0$ in some neighborhood of the origin



Stable




Stable




Stable

The origin is asymptotically stable if and only if $xf(x) < 0$ in some neighborhood of the origin



(a)

Asymptotically Stable



(b)

Globally Asymptotically Stable

Definition 3.2

Let the origin be an asymptotically stable equilibrium point of the system $\dot{x} = f(x)$, where f is a locally Lipschitz function defined over a domain $D \subset \mathbb{R}^n$ ($0 \in D$)


- The region of attraction (also called region of asymptotic stability, domain of attraction, or basin) is the set of all points x_0 in D such that the solution of

$$\dot{x} = f(x), \quad x(0) = x_0$$

is defined for all $t \geq 0$ and converges to the origin as t tends to infinity

- The origin is globally asymptotically stable if the region of attraction is the whole space $\mathbb{R}^n$

Example: Tunnel Diode Circuit



Linear Time-Invariant Systems

$$\dot{x} = Ax$$

$$x(t) = \exp(At)x(0)$$

$$P^{-1}AP = J = \text{block diag}[J_1, J_2, \dots, J_r]$$

$$J_i = \begin{bmatrix} \lambda_i & 1 & 0 & \dots & \dots & 0 \\ 0 & \lambda_i & 1 & 0 & \dots & 0 \\ \vdots & & \ddots & & & \vdots \\ \vdots & & & \ddots & & 0 \\ \vdots & & & & \ddots & 1 \\ 0 & \dots & \dots & \dots & 0 & \lambda_i \end{bmatrix}_{m \times m}$$

$$\exp(At) = P \exp(Jt) P^{-1} = \sum_{i=1}^r \sum_{k=1}^{m_i} t^{k-1} \exp(\lambda_i t) R_{ik}$$

m_i is the order of the Jordan block J_i

$$\operatorname{Re}[\lambda_i] < 0 \quad \forall i \quad \Leftrightarrow \quad \text{Asymptotically Stable}$$

$$\operatorname{Re}[\lambda_i] > 0 \quad \text{for some } i \quad \Rightarrow \quad \text{Unstable}$$

$$\operatorname{Re}[\lambda_i] \leq 0 \quad \forall i \quad \& \quad m_i > 1 \quad \text{for} \quad \operatorname{Re}[\lambda_i] = 0 \quad \Rightarrow \quad \text{Unstable}$$

$$\operatorname{Re}[\lambda_i] \leq 0 \quad \forall i \quad \& \quad m_i = 1 \quad \text{for} \quad \operatorname{Re}[\lambda_i] = 0 \quad \Rightarrow \quad \text{Stable}$$

If an $n \times n$ matrix A has a repeated eigenvalue λ_i of algebraic multiplicity q_i , then the Jordan blocks of λ_i have order one if and only if $\operatorname{rank}(A - \lambda_i I) = n - q_i$

Theorem 3.1

The equilibrium point $x = 0$ of $\dot{x} = Ax$ is stable if and only if all eigenvalues of A satisfy $\text{Re}[\lambda_i] \leq 0$ and for every eigenvalue with $\text{Re}[\lambda_i] = 0$ and algebraic multiplicity $q_i \geq 2$, $\text{rank}(A - \lambda_i I) = n - q_i$, where n is the dimension of x . The equilibrium point $x = 0$ is globally asymptotically stable if and only if all eigenvalues of A satisfy $\text{Re}[\lambda_i] < 0$

When all eigenvalues of A satisfy $\text{Re}[\lambda_i] < 0$, A is called a *Hurwitz matrix*

When the origin of a linear system is asymptotically stable, its solution satisfies the inequality

$$\|x(t)\| \leq k\|x(0)\|e^{-\lambda t}, \quad \forall t \geq 0, \quad k \geq 1, \quad \lambda > 0$$

Exponential Stability

Definition 3.3

The equilibrium point $x = 0$ of $\dot{x} = f(x)$ is exponentially stable if

$$\|x(t)\| \leq k\|x(0)\|e^{-\lambda t}, \quad \forall t \geq 0$$

$k \geq 1$, $\lambda > 0$, for all $\|x(0)\| < c$

It is globally exponentially stable if the inequality is satisfied for any initial state $x(0)$

Exponential Stability $\Rightarrow$ Asymptotic Stability

Example 3.2

$$\dot{x} = -x^3$$

The origin is asymptotically stable

$$x(t) = \frac{x(0)}{\sqrt{1 + 2tx^2(0)}}$$

$x(t)$ does not satisfy $|x(t)| \leq ke^{-\lambda t}|x(0)|$ because

$$|x(t)| \leq ke^{-\lambda t}|x(0)| \Rightarrow \frac{e^{2\lambda t}}{1 + 2tx^2(0)} \leq k^2$$

Impossible because $\lim_{t \rightarrow \infty} \frac{e^{2\lambda t}}{1 + 2tx^2(0)} = \infty$

Linearization

$$\dot{x} = f(x), \quad f(0) = 0$$

f is continuously differentiable over $D = \{\|x\| < r\}$

$$J(x) = \frac{\partial f}{\partial x}(x)$$

$$h(\sigma) = f(\sigma x) \quad \text{for } 0 \leq \sigma \leq 1, \quad h'(\sigma) = J(\sigma x)x$$

$$h(1) - h(0) = \int_0^1 h'(\sigma) d\sigma, \quad h(0) = f(0) = 0$$

$$f(x) = \int_0^1 J(\sigma x) d\sigma x$$

$$f(x) = \int_0^1 J(\sigma x) d\sigma x$$

Set $A = J(0)$ and add and subtract Ax

$$f(x) = [A + G(x)]x, \text{ where } G(x) = \int_0^1 [J(\sigma x) - J(0)] d\sigma$$

$$G(x) \rightarrow 0 \text{ as } x \rightarrow 0$$

This suggests that in a small neighborhood of the origin we can approximate the nonlinear system $\dot{x} = f(x)$ by its linearization about the origin $\dot{x} = Ax$

Theorem 3.2

- The origin is exponentially stable **if and only if** $\operatorname{Re}[\lambda_i] < 0$ for all eigenvalues of A
- The origin is unstable if $\operatorname{Re}[\lambda_i] > 0$ for some i

Linearization fails when $\operatorname{Re}[\lambda_i] \leq 0$ for all i , with $\operatorname{Re}[\lambda_i] = 0$ for some i

Example 3.3

$$\dot{x} = ax^3, \quad A = \left. \frac{\partial f}{\partial x} \right|_{x=0} = 3ax^2|_{x=0} = 0$$

Stable if $a = 0$; Asymp stable if $a < 0$; Unstable if $a > 0$

When $a < 0$, the origin is not exponentially stable

Lyapunov's Method

Let $V(x)$ be a continuously differentiable function defined in a domain $D \subset \mathbb{R}^n$; $0 \in D$. The derivative of V along the trajectories of $\dot{x} = f(x)$ is

$$\begin{aligned}\dot{V}(x) &= \sum_{i=1}^n \frac{\partial V}{\partial x_i} \dot{x}_i = \sum_{i=1}^n \frac{\partial V}{\partial x_i} f_i(x) \\ &= \left[\frac{\partial V}{\partial x_1}, \frac{\partial V}{\partial x_2}, \dots, \frac{\partial V}{\partial x_n} \right] \begin{bmatrix} f_1(x) \\ f_2(x) \\ \vdots \\ f_n(x) \end{bmatrix} \\ &= \frac{\partial V}{\partial x} f(x)\end{aligned}$$

If $\phi(t; x)$ is the solution of $\dot{x} = f(x)$ that starts at initial state x at time $t = 0$, then

$$\dot{V}(x) = \left. \frac{d}{dt} V(\phi(t; x)) \right|_{t=0}$$

If $\dot{V}(x)$ is negative, V will decrease along the solution of $\dot{x} = f(x)$

If $\dot{V}(x)$ is positive, V will increase along the solution of $\dot{x} = f(x)$

Lyapunov's Theorem (3.3)

- If there is $V(x)$ such that

$$V(0) = 0 \quad \text{and} \quad V(x) > 0, \quad \forall x \in D \text{ with } x \neq 0$$

$$\dot{V}(x) \leq 0, \quad \forall x \in D$$

then the origin is a stable

- Moreover, if

$$\dot{V}(x) < 0, \quad \forall x \in D \text{ with } x \neq 0$$


then the origin is asymptotically stable

- Furthermore, if $V(x) > 0, \forall x \neq 0$,

$$\|x\| \rightarrow \infty \Rightarrow V(x) \rightarrow \infty$$

and $\dot{V}(x) < 0, \forall x \neq 0$, then the origin is globally asymptotically stable

Proof



$$0 < r \leq \varepsilon, \quad B_r = \{\|x\| \leq r\}$$

$$\alpha = \min_{\|x\|=r} V(x) > 0$$

$$0 < \beta < \alpha$$

$$\Omega_\beta = \{x \in B_r \mid V(x) \leq \beta\}$$

$$\|x\| \leq \delta \Rightarrow V(x) < \beta$$

Solutions starting in Ω_β stay in Ω_β because $\dot{V}(x) \leq 0$ in Ω_β

$$x(0) \in B_\delta \Rightarrow x(0) \in \Omega_\beta \Rightarrow x(t) \in \Omega_\beta \Rightarrow x(t) \in B_r$$

$$\|x(0)\| < \delta \Rightarrow \|x(t)\| < r \leq \varepsilon, \quad \forall t \geq 0$$

$\Rightarrow$ The origin is stable

Now suppose $\dot{V}(x) < 0 \quad \forall x \in D, x \neq 0$. $V(x(t))$ is monotonically decreasing and $V(x(t)) \geq 0$

$$\lim_{t \rightarrow \infty} V(x(t)) = c \geq 0 \quad \text{Show that } c = 0$$

Suppose $c > 0$. By continuity of $V(x)$, there is $d > 0$ such that $B_d \subset \Omega_c$. Then, $x(t)$ lies outside B_d for all $t \geq 0$

$$\gamma = - \max_{d \leq \|x\| \leq r} \dot{V}(x)$$

$$V(x(t)) = V(x(0)) + \int_0^t \dot{V}(x(\tau)) d\tau \leq V(x(0)) - \gamma t$$

This inequality contradicts the assumption $c > 0$

$\Rightarrow$ The origin is asymptotically stable

The condition $\|x\| \rightarrow \infty \Rightarrow V(x) \rightarrow \infty$ implies that the set $\Omega_c = \{x \in R^n \mid V(x) \leq c\}$ is compact for every $c > 0$. This is so because for any $c > 0$, there is $r > 0$ such that $V(x) > c$ whenever $\|x\| > r$. Thus, $\Omega_c \subset B_r$. All solutions starting Ω_c will converge to the origin. For any point $p \in R^n$, choosing $c = V(p)$ ensures that $p \in \Omega_c$

$\Rightarrow$ The origin is globally asymptotically stable


Terminology

$V(0) = 0, V(x) \geq 0 \text{ for } x \neq 0$	Positive semidefinite
$V(0) = 0, V(x) > 0 \text{ for } x \neq 0$	Positive definite
$V(0) = 0, V(x) \leq 0 \text{ for } x \neq 0$	Negative semidefinite
$V(0) = 0, V(x) < 0 \text{ for } x \neq 0$	Negative definite
$\ x\ \rightarrow \infty \Rightarrow V(x) \rightarrow \infty$	Radially unbounded

Lyapunov' Theorem

The origin is stable if there is a continuously differentiable positive definite function $V(x)$ so that $\dot{V}(x)$ is negative semidefinite, and it is asymptotically stable if $\dot{V}(x)$ is negative definite. It is globally asymptotically stable if the conditions for asymptotic stability hold globally and $V(x)$ is radially unbounded

A continuously differentiable function $V(x)$ satisfying the conditions for stability is called a *Lyapunov function*. The surface $V(x) = c$, for some $c > 0$, is called a *Lyapunov surface* or a *level surface*




Why do we need the radial unboundedness condition to show global asymptotic stability?

It ensures that $\Omega_c = \{V(x) \leq c\}$ is bounded for every $c > 0$. Without it Ω_c might not be bounded for large c

Example

$$V(x) = \frac{x_1^2}{1 + x_1^2} + x_2^2$$



Example: Pendulum equation without friction

$$\dot{x}_1 = x_2, \quad \dot{x}_2 = -a \sin x_1$$

$$V(x) = a(1 - \cos x_1) + \frac{1}{2}x_2^2$$

$V(0) = 0$ and $V(x)$ is positive definite over the domain $-2\pi < x_1 < 2\pi$

$$\dot{V}(x) = a\dot{x}_1 \sin x_1 + x_2\dot{x}_2 = ax_2 \sin x_1 - ax_2 \sin x_1 = 0$$

The origin is stable

Since $\dot{V}(x) \equiv 0$, the origin is not asymptotically stable

Example: Pendulum equation with friction

$$\dot{x}_1 = x_2, \quad \dot{x}_2 = -a \sin x_1 - bx_2$$

$$V(x) = a(1 - \cos x_1) + \frac{1}{2}x_2^2$$

$$\dot{V}(x) = a\dot{x}_1 \sin x_1 + x_2\dot{x}_2 = -bx_2^2$$

The origin is stable

$\dot{V}(x)$ is not negative definite because $\dot{V}(x) = 0$ for $x_2 = 0$ irrespective of the value of x_1

The conditions of Lyapunov's theorem are only sufficient. Failure of a Lyapunov function candidate to satisfy the conditions for stability or asymptotic stability does not mean that the equilibrium point is not stable or asymptotically stable. It only means that such stability property cannot be established by using this Lyapunov function candidate

Try

$$\begin{aligned} V(x) &= \frac{1}{2}x^T P x + a(1 - \cos x_1) \\ &= \frac{1}{2} \begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} p_{11} & p_{12} \\ p_{12} & p_{22} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + a(1 - \cos x_1) \end{aligned}$$

$$p_{11} > 0, \quad p_{11}p_{22} - p_{12}^2 > 0$$

$$\begin{aligned}
 \dot{V}(x) &= (p_{11}x_1 + p_{12}x_2 + a \sin x_1) x_2 \\
 &\quad + (p_{12}x_1 + p_{22}x_2) (-a \sin x_1 - bx_2) \\
 &= a(1 - p_{22})x_2 \sin x_1 - ap_{12}x_1 \sin x_1 \\
 &\quad + (p_{11} - p_{12}b) x_1x_2 + (p_{12} - p_{22}b) x_2^2
 \end{aligned}$$

$$p_{22} = 1, \quad p_{11} = bp_{12} \Rightarrow 0 < p_{12} < b, \quad \text{Take } p_{12} = b/2$$

$$\dot{V}(x) = -\frac{1}{2}abx_1 \sin x_1 - \frac{1}{2}bx_2^2$$

$$D = \{|x_1| < \pi\}$$

$V(x)$ is positive definite and $\dot{V}(x)$ is negative definite over D .
The origin is asymptotically stable

Variable Gradient Method

$$\dot{V}(x) = \frac{\partial V}{\partial x} f(x) = g^T(x) f(x)$$

$$g(x) = \nabla V = (\partial V / \partial x)^T$$

Choose $g(x)$ as the gradient of a positive definite function $V(x)$ that would make $\dot{V}(x)$ negative definite

$g(x)$ is the gradient of a scalar function if and only if

$$\frac{\partial g_i}{\partial x_j} = \frac{\partial g_j}{\partial x_i}, \quad \forall i, j = 1, \dots, n$$

Choose $g(x)$ such that $g^T(x)f(x)$ is negative definite

Compute the integral

$$V(x) = \int_0^x g^T(y) \, dy = \int_0^x \sum_{i=1}^n g_i(y) \, dy_i$$

over any path joining the origin to x ; for example

$$\begin{aligned} V(x) = & \int_0^{x_1} g_1(y_1, 0, \dots, 0) \, dy_1 + \int_0^{x_2} g_2(x_1, y_2, 0, \dots, 0) \, dy_2 \\ & + \dots + \int_0^{x_n} g_n(x_1, x_2, \dots, x_{n-1}, y_n) \, dy_n \end{aligned}$$

Leave some parameters of $g(x)$ undetermined and choose them to make $V(x)$ positive definite

Example 3.7 (Read)

$$\dot{x}_1 = x_2, \quad \dot{x}_2 = -h(x_1) - ax_2$$

$a > 0$, $h(\cdot)$ is locally Lipschitz,

$$h(0) = 0; \quad yh(y) > 0 \quad \forall y \neq 0, \quad y \in (-b, c), \quad b > 0, \quad c > 0$$

$$\frac{\partial g_1}{\partial x_2} = \frac{\partial g_2}{\partial x_1}$$

$$\dot{V}(x) = g_1(x)x_2 - g_2(x)[h(x_1) + ax_2] < 0, \quad \text{for } x \neq 0$$

$$V(x) = \int_0^x g^T(y) dy > 0, \quad \text{for } x \neq 0$$

$$\text{Try} \quad g(x) = \begin{bmatrix} \phi_1(x_1) + \psi_1(x_2) \\ \phi_2(x_1) + \psi_2(x_2) \end{bmatrix}$$

To satisfy the symmetry requirement, we must have

$$\frac{\partial \psi_1}{\partial x_2} = \frac{\partial \phi_2}{\partial x_1}$$

$$\psi_1(x_2) = \gamma x_2 \quad \text{and} \quad \phi_2(x_1) = \gamma x_1$$

$$\begin{aligned} \dot{V}(x) = & -\gamma x_1 h(x_1) - a x_2 \psi_2(x_2) + \gamma x_2^2 \\ & + x_2 \phi_1(x_1) - a \gamma x_1 x_2 - \psi_2(x_2) h(x_1) \end{aligned}$$

To cancel the cross-product terms, take

$$\psi_2(x_2) = \delta x_2 \quad \text{and} \quad \phi_1(x_1) = a\gamma x_1 + \delta h(x_1)$$

$$g(x) = \begin{bmatrix} a\gamma x_1 + \delta h(x_1) + \gamma x_2 \\ \gamma x_1 + \delta x_2 \end{bmatrix}$$

$$\begin{aligned} V(x) &= \int_0^{x_1} [a\gamma y_1 + \delta h(y_1)] dy_1 + \int_0^{x_2} (\gamma x_1 + \delta y_2) dy_2 \\ &= \frac{1}{2}a\gamma x_1^2 + \delta \int_0^{x_1} h(y) dy + \gamma x_1 x_2 + \frac{1}{2}\delta x_2^2 \\ &= \frac{1}{2}x^T P x + \delta \int_0^{x_1} h(y) dy, \quad P = \begin{bmatrix} a\gamma & \gamma \\ \gamma & \delta \end{bmatrix} \end{aligned}$$

$$V(x) = \frac{1}{2}x^T Px + \delta \int_0^{x_1} h(y) dy, \quad P = \begin{bmatrix} a\gamma & \gamma \\ \gamma & \delta \end{bmatrix}$$

$$\dot{V}(x) = -\gamma x_1 h(x_1) - (a\delta - \gamma)x_2^2$$

Choose $\delta > 0$ and $0 < \gamma < a\delta$

If $yh(y) > 0$ holds for all $y \neq 0$, the conditions of Lyapunov's theorem hold globally and $V(x)$ is radially unbounded